



Europäisches
Patentamt

European
Patent Office

Office européen
des brevets

REC'D: 17 SEP 2004

WIPO

PCT

Bescheinigung

Certificate

Attestation

Die angehefteten Unterla-
gen stimmen mit der
ursprünglich eingereichten
Fassung der auf dem näch-
sten Blatt bezeichneten
europäischen Patentanmel-
dung überein.

The attached documents
are exact copies of the
European patent application
described on the following
page, as originally filed.

Les documents fixés à
cette attestation sont
conformes à la version
initialement déposée de
la demande de brevet
européen spécifiée à la
page suivante.

Patentanmeldung Nr. Patent application No. Demande de brevet n°

03077451.7

03077451.7

Der Präsident des Europäischen Patentamts;
Im Auftrag

For the President of the European Patent Office

Le Président de l'Office européen des brevets
p.o.

R C van Dijk

**PRIORITY
DOCUMENT**

SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH RULE 17.1(a) OR (b)



Anmeldung Nr:
Application no.: 03077451.7
Demande no:

Anmeldetag:
Date of filing: 05.08.03
Date de dépôt:

Anmelder/Applicant(s)/Demandeur(s):

Fuji Photo Film B.V.
Oudenstaart 1
5047 TK Tilburg
PAYS-BAS

Bezeichnung der Erfindung/Title of the invention/Titre de l'invention:
(Falls die Bezeichnung der Erfindung nicht angegeben ist, siehe Beschreibung.
If no title is shown please refer to the description.
Si aucun titre n'est indiqué se referer à la description.)

Use of gelatin-like proteins as stabiliser

In Anspruch genommene Priorität(en) / Priority(ies) claimed /Priorité(s)
revendiquée(s)
Staat/Tag/Aktenzeichen/State/Date/File no./Pays/Date/Numéro de dépôt:

Internationale Patentklassifikation/International Patent Classification/
Classification internationale des brevets:

A61K47/00

Am Anmeldetag benannte Vertragstaaten/Contracting states designated at date of
filing/Etats contractants désignées lors du dépôt:

AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HU IE IT LU MC NL
PT RO SE SI SK TR LI

Use of gelatin-like proteins as stabiliser**05. 08. 2003**

(87)

FIELD OF THE INVENTION

- 5 The invention relates to the use of gelatin-like proteins - or polypeptides - as stabilisers in lyophilized biological or pharmaceutical compositions.

BACKGROUND OF THE INVENTION

- 10 A well-established application of gelatin is the use as stabilizer for physiologically active substances in lyophilized biological or pharmaceutical compositions.

- Lyophilization or freeze drying of physiologically active substances is generally done in the presence of a stabiliser and a disaccharide. Freeze drying compositions and - processes are empirically determined for different types of physiologically active substances, as described by D. Greiff in Developments in Biological Standardization
15 (1992), 7 Biol. Prod. Freeze Drying Formulation), 85-92. The stability of the lyophilized composition depends on several factors like the nature of the physiologically active substance, and water content and glass transition temperature (Tg) of the freeze-dried composition. Vaccines are examples of pharmaceutical compounds stored as freeze-dried compositions.

- 20 Vaccines are used amongst others in development countries where the sometimes severe storage conditions for vaccines can be difficult to maintain. Stability of lyophilized vaccines is a major concern, and the World Health Organisation issues strict rules for storage of such compositions.

- 25 Physiologically active substances are for example vaccines, (therapeutic) proteins, enzymes, (monoclonal) antibodies and the like. Gelatin is a preferred stabiliser because of its known low immunogenicity. Care should be taken that the gelatin solution is made sterile, pyrogen and antigen free.

A disadvantage of the presently used gelatin is the possibility of immediate hypersensitivity, which can occur upon application of the presently used gelatin derivatives, known as anaphylactic shock.

5 Another disadvantage of the commercially used gelatin derivatives is the fact that the gelatin used is isolated from animal sources such as animal bone and hide, in particular it is derived from bovine sources. Disadvantages of this material are the presence of impurities and the fact that the nature of the composition is not clearly defined and thus not reproducible. This may impose additional screening to ensure that the derivatisation
10 process results in a product with the desired properties and may require careful purification steps. An additional problem nowadays, especially in relation to gelatin isolated from bovine sources, is the risk of contamination of the gelatin with factors responsible for the occurrence of Bovine Spongiform Encephalitis (BSE). For this reason the use of gelatin in pharmaceutical compositions may be prohibited.

15

WO 01/34801 A2 describes generally the use of recombinant gelatins as vaccine stabiliser to avoid the obvious problems associated with the use of natural gelatin. However, it is silent with respect to further advantages, which can be achieved by specifically designed recombinant structures.

20

EP 0,781,779 A2 describes the use of a gelatin of not more than 20 kiloDalton (kDa) that is hydrolyzed specifically by collagenase to render it non-antigenic. US 4,147,772 describes the use of hydrolyzed gelatin of about 3 kDa as a nongelling matrix with little antigenicity.

25

US 4,273,762 describes an attempt to reduce the lyophilization time of vaccines which have partly hydrolyzed gelatin as stabiliser.

SUMMARY OF THE INVENTION

30 It is an object of the invention to provide improved stabilisers for lyophilized compositions comprising physiologically active substances.

It is also an object of the invention to provide lyophilized compositions comprising the improved stabilisers, said compositions having an improved stability.

It is a further object of the invention to reduce the lyophilizing time for compositions comprising physiologically active substances with the improved stabiliser.

Surprisingly it was found that these objectives were met by using as a stabilizer a recombinant or synthetic polypeptide comprising at least one stretch of 10 or more consecutive repeats of Gly-Xaa-Yaa triplets and in which at least 20% of the amino acids are present in the form of consecutive Gly-Xaa-Yaa triplets and said recombinant or synthetic polypeptide having a calculated glass transition temperature of higher than about 180 degrees Celsius, as calculated by formula 8 and 9 of Matveev as published in Food Hydrocolloids Vol. 11 no.2 pp. 125-133, 1997. A peptide with these characteristics is hereinafter referred to as "recombinant" or "synthetic collagen-like peptide (or polypeptide)" or "recombinant" or "synthetic gelatin-like peptide (or polypeptide)", depending on the method of its production (i.e. by recombinant expression or by chemical synthesis).

It was also found that the lyophilization process can be optimized significantly when the recombinant polypeptide of the invention has no helical structure.

DESCRIPTION OF THE INVENTION

According to the invention a lyophilized composition is provided comprising as a stabilizer a recombinant or synthetic polypeptide with a calculated glass transition temperature that is higher than about 180 degrees Celsius, comprising at least one stretch of 10 or more consecutive repeats of Gly-Xaa-Yaa triplets and in which at least 20% of the amino acids are present in the form of consecutive Gly-Xaa-Yaa triplets.

The measured glass transition temperature of the composition should also be significantly higher, preferably at least about 5 degrees, more preferably at least about 10 degrees and most preferably 20 degrees Celsius higher, than the measured glass transition temperature of a control composition, which comprises native collagen

peptides. "Native collagen" as used herein refers to collagen peptides or polypeptides which were not selected or synthesized to have a high glass transition temperature. In general, native collagen peptides have a calculated Tg of about 170 degrees Celsius or less.

5

It is noted, that when the Tg of a mixture, composed of a gelatin-like peptide and one or more other compounds, is measured, the measured Tg of the composition may be significantly different from the measured Tg of the substantially pure gelatin-like peptide. For example, the measured Tg of a composition comprising a gelatin-like peptide and sucrose may be significantly lower than the measured Tg of the pure gelatin-like peptide.

10

Pharmaceutical formulations that are introduced into the bloodstream contain proteins as, for example, a stabiliser, as a drug carrier or as an osmotic colloid. It is long recognized in the art that gelatins are preferred for their low immunogenicity. It is also recognized in the art that recombinant gelatins can advantageously replace gelatins from natural sources to avoid introduction of non-gelatin material. Recently the occurrence of BSE has been a source of concern and a reason to avoid the use of gelatin from natural sources.

15
20

Although the use of recombinant gelatins is described for the obvious reasons, and it is suggested that recombinant structures can be optimised there are no teachings as to what such optimisation might comprise.

25

In our studies on collagen properties we found to our surprise that, although collagen has a repetitive amino acid triplet structure Gly-Xaa-Yaa, wherein a majority of the triplets contain a proline, the glass transition temperature (or Tg) is not uniformly divided over the molecule, and sequences can be selected that have a higher Tg than the average (native) collagen.

30

The importance of the glass transition temperature is well known in the art of freeze drying or lyophilizing of formulations containing physiologically active substances,

like vaccines. In lyophilized formulations one strives for high glass transition temperature. In "Long-Term Stabilization of Biologicals" (Biotechnology vol.12 12 march 1994) F. Franks addresses the importance of high glass transition temperatures in the preservation of biological materials by freeze drying and the desire to further
5 improve the shelf life of such materials. In the formulations for freeze drying, gelatin serves to protect the physiologically active substance whereby the presence of water molecules bound to polar groups of the amino acid residues is thought to be of importance. Residual moisture plays an important role in the shelf life of vaccines. Increased residual moisture levels decrease the glass transition temperature of a
10 lyophilized gelatin/disaccharide composition significantly, resulting in reduced shelf life.

There are many publications on this subject, for example by Phillips et. al in cryobiology 18, 414-419 (1981) or US 801,856. Vaccines like MMR (Mumps Measles
15 Rubella) have in current formulations a critical Tg, which lies around 47 degrees Celsius under dry conditions but rapidly decreases towards room temperature when small amounts of moisture enter the material. Within one week at 37 degrees Celsius a loss in potency of 50% is reported by M.K. Lala in Indian Pediatrics 2003; 40:311-319 Increasing the Tg even by a few degrees can have a tremendous effect on the shelf life
20 of these vaccines. The problem of a reduced stability of the physiologically active formulations, which are stabilized with gelatins, was solved by the present invention, which is based on the use of new recombinant or synthetic gelatins with an increased Tg in combination with a certain similarity with natural human gelatin amino acid sequences to prevent the occurrence of unwanted immune responses.

25

A recombinant or synthetic gelatin-like polypeptide according to the invention is preferably a sequence identical to or highly homologous to a native human collagen sequence. To select such an amino acid sequence from a native sequence, "moving Tg averages" (as defined below) are calculated. A sequence is then selected which has a
30 calculated average glass transition temperature of about 10 degrees Celsius higher than the calculated average collagen glass transition temperature of the native starting sequence, preferably about 20 degrees higher, more preferably about 30 degrees higher,

- even more preferably about 40 degrees higher. This value may differ somewhat between different types of collagen and depend on the presence of propeptides, telopeptides or signal peptides. The average calculated glass transition temperature of native collagen is about 170 degrees Celsius, so that a polypeptide according the invention has a T_g higher than about 180 degrees, preferably higher than about 190 degrees, more preferably higher than about 200 degrees. "About" as used herein refers to a temperature range of 1-4 degrees higher and/or lower than the specified temperature.
- T_g increases of less than 10 degrees are also considered, but the effect in the eventual formulation in which disaccharides are present may be reduced to a less significant level.

- The calculation method of the glass transition temperature was published by Y. Matveev et. al. in Food Hydrocolloids Vol. 11 no. 2 pp. 125-133, 1997. Equations 8 and 9 were used for the actual calculations:

$$(8) T_g^{-1} = \sum_{i=1}^{20} \phi_i T_{gi}^{-1} \quad \text{wherein} \quad (9) \phi_i = n_i \Delta V_i / \sum_{i=1}^{20} n_i \Delta V_i$$

- wherein the summations i=1 to 20 are the summations of the values for the partial values of T_g and ΔV of the separate amino acids given below (V is a measure for the vd Waals volume, as described in Matveev et al. (supra)):

| No. | Amino Acid | T _{gi} (Kelvin) | ΔV _i |
|-----|------------|--------------------------|-----------------|
| 1 | gly | 599 | 47.3 |
| 2 | ala | 621 | 64.4 |
| 3 | val | 931 | 98.6 |
| 4 | leu | 400 | 115.7 |
| 5 | ile | 400 | 115.7 |
| 6 | phe | 528 | 139.9 |
| 7 | pro | 423 | 88.0 |

| | | | |
|----|--------|-----|-------|
| 8 | trp | 544 | 196.9 |
| 9 | ser | 311 | 66.1 |
| 10 | thr | 321 | 88.9 |
| 11 | met | 362 | 120.6 |
| 12 | asn | 232 | 94.6 |
| 13 | gln | 312 | 111.7 |
| 14 | cys-SH | 418 | 82.2 |
| 15 | asp | 672 | 80.1 |
| 16 | glu | 487 | 97.2 |
| 17 | tyr | 573 | 136.9 |
| 18 | his | 488 | 118.9 |
| 19 | lys | 258 | 118.1 |
| 20 | arg | 410 | 138.4 |

The model does not appear to take the presence of hydroxyproline into account. However, the correlation with measured values which are presented in the paper of Matveev et al. give a very good correlation between calculated and measured values of gelatin.

For selecting appropriate recombinant or synthetic collagen-like peptides a starting point is for example human Col1A1 (SEQ ID NO: 1), which has a Tg of 163 degrees Celsius calculated from entire sequence.

SEQ ID NO: 1 (human Col1A1):

MFSFVDLRLLLLLAATALLTHGQEEGQVEGQDEDIPPITCVQNGRLRYHDRVW
 KPEPCRICVCDNGKVLCDDEVICDETKNCPGAEVPEGECCPVCPDGSESPTDQET
 TGVEGPKGDTGPRGPRGPAGPPGRDGIPGQPGLPGPPGPPGPPGPPGLGGNFAP
 QLSYGYDEKSTGGISVPGPMGPSGPRGLPGPPGAPGPQGFGPPGEPGEPGASG
 PMGPRGPPGPPGKNGDDGEAGKPGRPGERGPPGPQGARGLPGTAGLPGMKGH
 RGFSGLDGAKGDAGPAGPKGEPGSPGENGAPQMGPRLPGERGRPGAPGPA
 GARGNDGATGAAGPPGPTGPAGPPGFPGAVGAKGEAGPQGPRGSEGPQGVRG

EPGPPGPAGAAGPAGNPGADGQPGAKGANGAPGLAGAPGFPGARGPSGPQGP
 GGPPGPKGNSGEPGAPGSKGDTGAKGEPGPVGVQGPPGPAGEEGKRGARGE
 GPTGLPGPPGERGGPGSRGFPADGVAGPKGPAGERGSPGPAGPKGSPGEAGR
 PGEAGLPGAKGLTGSPGSPGPDGKTGPPGPAGQDGRPGPPGPPGARGQAGVM
 5 GFGPKGAAGEPGKAGERGVPGPPGAVGPAGKDGEAGAQQPPGPAGPAGERG
 EQGPAGSPGFQGLPGPAGPPGEAGKPGEQGVPGDLGAPGPSGARGERGFPGER
 GVQGPPGPAGPRGANGAPGNDGAKGDAGAPGAPGSQGAPGLQGMPPGERGAA
 GLPGPKGDRGDAGPKGADGSPGKDGVRLTGPIGPPGPAGAPGDKGESGPSGP
 AGPTGARGAPGDRGEPGPPGPAGFAGPPGADGQPGAKGEPGDAGAKGDAGPP
 10 GPAGPAGPPGPIGNVGAPGAKGARGGAGPPGATGFPGAAGRVPVGGPSGNAGP
 PGPPGPAGKEGGKGRGETGPAGRPGEVGPPGPPGPAGEKGSPPGADGPAGAPG
 TPGPQGLIAGQRGVVGLPGQRGERGFPLGPSGEPGKQGPSGASGERGPPGPM
 GPPGLAGPPGESGREGAPGAEGSPGRDGSPGAKGDRGETGPAGPPGAPGAPGA
 PGPVGPAGKSGDRGETGPAGPAGPVGPAGARGPAGPQGPRGDKGETGEQGDR
 15 GIKGHRGFSGGLQGPPGPPGSPGEQGPSGASGPAGPRGPPGSAGAPGKDGLNGLP
 GPIGPPGPRGRTGDAGPVGPPGPPGPPGPPGPPSAGFDFSLPQPPQEK AHDGGR
 YYRADDANVVRDRDLEVDTTLSLSQQIENIRSPEGSRKNPARTCRDLKMCHS
 DWKSGEYWIDPNQGCNLDAIKVFCNMETGETCVYPTQPSVAQKNWYISKNP
 DKRHVWFGESMTDGFQFEYGGQGSDDPADVAIQLTFLRLMSTEASQNITYHCK
 20 NSVAYMDQQTGNLKKALLLKGSNEIEIRAEGNSRFTYSVTVDGCTSHTGAWG
 KTVIEYKTT KTSRLPIIDVAPLDVGAPDQEFGFDVGPVCFL

This Col1A1 sequence still includes the signal sequence (amino acids 1-22) and the
 amino terminal propeptides (amino acids 23-161 and 1219-1464). The helical collagen
 25 sequence is present from amino acid 162 to amino acid 1218. Using a spreadsheet the
 moving average over a number of amino acids could easily be calculated and displayed.
 Figures 1 to 4 show the result for a moving average of resp. 18, 27, 54 and 81 amino
 acids. A "moving Tg average" of, for example $n=54$, means that first the average Tg of
 the first to the 54th amino acid is calculated, then of the 2nd to 55th amino acid, then from
 30 the 3rd to the 56th and so on. These values are then plotted as in fig. 3, the first datapoint
 being plotted at the 54th amino acid. The amino acid regions which have a calculated Tg
 higher than the average calculated Tg of this native collagen (i.e. the average calculated

Tg of the complete sequence) can now be identified. It is remarkable that smaller polypeptides allow selection of regions with higher Tg. Calculating a moving average of 54 amino acids allows selection of polypeptide sequences with increased Tg of up to about 200 degrees C. For example a sequence from amino acid 1034 to 1087 of SEQ ID NO: 1 results in a calculated Tg of 208 degrees Celsius. This polypeptide has, thus, a calculated Tg which is 45 degrees Celsius higher than the calculated Tg of the native sequence, which is 163 degrees Celsius calculated for entire sequence. When expressed as such this yields a gelatin-like polypeptide of about 5,000 Dalton. A sequence of about 500 amino acids can be selected from about amino acid 600 to about amino acid 1100 of SEQ ID NO: 1, that still has an average Tg of about 178 degrees Celsius and a molecular weight of about 40,000 to 50,000 Dalton. From about amino acid 590 to 750 of SEQ ID NO: 1 a polypeptide with an average Tg of higher than 180 degrees Celsius can be selected that has a molecular weight of up to about 10,000 to 13,000 Dalton. Polypeptide regions with the desired average Tg such as described here above can be easily calculated also from other collagen sequences, such as Col 1A-2, Col 2A-1, Col 3A-1 and so on. Such collagen sequences are readily available in the art.

When desired, repetitive sequences of these sequences can be expressed to obtain larger molecular weights. Conventional hydrolysed gelatins with a weight of about 3,000 to 15,000 Dalton are applied, preferably between 5,000 and 10,000 Dalton and more preferably between 6,000 and 8,000 Dalton. When desired also larger molecular weights can be obtained by the invention giving a specific advantage for the achievable Tg. Thus in one embodiment the gelatin-like polypeptide has a preferred molecular weight between 3,000 and 15,000 Dalton, more preferably between 5,000 and 10,000, even more preferably between 6,000 and 8,000 Dalton. In another embodiment the gelatin-like polypeptide has a molecular weight between 3,000 and 80,000 Dalton, preferably between 5,000 and 60,000 Dalton, most preferably between 10,000 and 40,000 Dalton.

It was attempted to correlate the Tg of a polypeptide fragment to its structural details. Some correlation was found with the alanine content, as shown in figure 5. Although for a moving average of 54 amino acids many of the areas with higher Tg coincide with

elevated alanine levels, this correlation is not valid for all regions with a Tg higher than average. Still, with a moving average of 54 amino acids it is likely that a region with higher Tg is found when the polypeptide of 54 amino acids has an alanine content of more than about 1 alanine per 10 amino acids. The presence of bulky amino acid residues can have a negative effect on the Tg of a polypeptide. A correlation was made between the presence of leucine and isoleucine and the Tg over a moving average of 54 amino acids (fig. 6). In many areas with high Tg, but not all, the concentration of these bulky amino acid residues is low, or they are absent. Bringing valine in the correlation makes it worse, suggesting that valine has less effect on the bulkiness. Considering the sizes of the side chains of the abundantly present prolines it is imaginable that leucine and isoleucine contribute more to the bulkiness than valine. Further, it is desirable that the amount of polar amino acid residues is more than 5% and more preferably more than 7% but less than 15% so that enough water molecules can be bound to protect the lyophilized physiologically active substance.

Gelatin-like recombinant or synthetic polypeptides according to the invention are preferably identical or essentially similar to natural human collagen amino acid sequences, but also non-human sequences (such as rat, rabbit, mouse etc.) can be used, or sequences can be designed that do not occur naturally. The term "essentially similar" means that two peptide sequences, when optimally aligned, such as by the programs GAP or BESTFIT using default parameters, share at least 80 percent sequence identity, preferably at least 90 percent sequence identity, more preferably at least 95 percent sequence identity or more (e.g., 99 or 100 percent sequence identity). GAP uses the Needleman and Wunsch global alignment algorithm to align two sequences over their entire length, maximizing the number of matches and minimizes the number of gaps. Generally, the GAP default parameters are used, with a gap creation penalty = 50 (nucleotides) / 8 (proteins) and gap extension penalty = 3 (nucleotides) / 2 (proteins).

Such sequences would preferably have a high alanine content of more than 10 alanine residues per 100 amino acids, preferably more than 12 per 100 amino acids, more preferably more than 14 per 100 amino acids. Such a designed structure contains polar

amino acid residues comparable to natural gelatins. The incorporation of bulky amino acids is to be avoided.

A natural gelatin molecule in its primary amino acid sequence basically consists of repeats of Gly-Xaa-Yaa triplets, thus approximately one third of the total number of amino acids is a glycine. The molecular weight of gelatin is typically large, values of the molecular weight vary from 10,000 to 300,000 daltons. The main fraction of natural gelatin molecules has a molecular weight around 90,000 daltons. The average molecular weight is higher than 90,000 daltons.

Furthermore, characteristic for gelatin is the unusual high content of proline residues. Even more characteristic is that in natural gelatin a number of the proline residues is hydroxylated. Most prominent site of hydroxylation is the 4-position resulting in the presence in the gelatin molecule of the unusual amino acid 4-hydroxyproline. In a triplet 4-hydroxyproline is always found in the Yaa position. Very few proline residues are hydroxylated at the 3 position. In contrast with 4-hydroxyproline, 3-hydroxyproline is always found at the carboxyl side of a glycine residue, thus in the Xaa position in a triplet. Different enzymes are responsible for the formation of 3- or 4-hydroxyproline.

Based on known amino acid compositions, it is estimated that in a gelatin molecule derived from a mammal, approximately 22 % of the amino acids are a proline or a hydroxyproline residue. However lower contents of proline and hydroxyproline are found in fish, in particular cold water fish. A rough estimate is that proline and hydroxyproline residues are present in approximately equal amounts, thus in a gelatin molecule derived from a mammal approximately 11 % of the amino acids are prolines and approximately 11 % are hydroxyprolines. As substantially all hydroxyproline is found in the Yaa position, it is estimated that approximately one third of all triplets in a gelatin molecule comprise a hydroxyproline. The presence of the hydroxyproline residues is responsible for the fact that a gelatin molecule in its secondary structure can adopt a helical conformation.

Furthermore, another amino acid present in natural gelatin that is found in very few other proteins is 5-hydroxylysine. Lysine residues modified in this way are always found in the Yaa position in a triplet.

5 A predominant feature of gelatins is the presence of Gly-Xaa-Yaa triplets. Such triplets are also present in the gelatin-like proteins of this invention. It is however possible to design a protein in which Gly-Xaa-Yaa triplets or stretches of Gly-Xaa-Yaa triplets are separated by one or more amino acids without significantly altering the gelatin-like character of the protein. Such gelatin-like proteins are comprised by the definition of
10 gelatin-like protein of this invention.

The gelatin-like proteins for use according to the invention can be produced by recombinant methods as disclosed in EP-A-0926543 and EP-A-1014176. For enablement of the production and purification of gelatin-like proteins that can be
15 suitably used in composition according to the invention specific reference is made to the examples in EP-A-0926543 and EP-A-1014176. Thus the gelatin-like proteins can be produced by expression of nucleic acid sequence encoding such polypeptide by a suitable microorganism. The process can suitably be carried out with a fungal cell or a yeast cell. Suitably the host cell is a high expression host cell like Hansenula,
20 Trichoderma, Aspergillus, Penicillium, Neurospora or Pichia. Fungal and yeast cells are preferred to bacteria as they are less susceptible to improper expression of repetitive sequences. Most preferably the host will not have a high level of proteases that attack the collagen structure expressed. In this respect Pichia offers an example of a very suitable expression system. As disclosed in EP-A-0926543 and EP-A-1014176
25 specifically Pichia pastoris is used as expression system. In one embodiment the microorganism is also transformed to include a gene for expression of prolyl-4-hydroxylase. In another embodiment the microorganism is free of active post-translational processing mechanism such as in particular hydroxylation of proline.

30 The selection of a suitable host cell from known industrial enzyme producing fungal host cells specifically yeast cells on the basis of the required parameters described herein rendering the host cell suitable for expression of recombinant gelatin-like proteins

suitable in compositions according to the invention in combination with knowledge regarding the host cells and the sequence to be expressed will be possible by a person skilled in the art.

5 With respect to the design of gelatin-like proteins for use in the invention, several properties of the proteins are addressed. For instance it can be made sure specific amino acids, such as bulky amino acids like leucine or isoleucine which lower the average T_g, will not occur in the protein or only occur infrequently. Otherwise, as discussed above in particular with respect to alanine or polar amino acids, it can be advantageous to
10 introduce a definite number of a specific amino acid in the gelatin-like protein. Yet further the iso-electric point (IEP) can be tuned by the composition of acidic and basic amino acid residues in the gelatin-like proteins.

In one embodiment the composition according to the invention comprises a gelatin-like
15 protein which is homodisperse in nature. Homodisperse means of constant composition and molecular weight. Variations in composition that can occur due to the recombinant production process are allowed. In terms of molecular weight a useful definition of homodispersity would be that at least 90% of the total amount of gelatin-like protein in the composition has a molecular weight that lies within a range of plus or minus 10%
20 around a selected molecular weight. In another embodiment the composition according to the invention comprises two or more gelatin-like proteins each being homodisperse in nature but with different molecular weights (i.e. a bimodal molecular weight distribution). This prevents crystallization during the freeze drying process or during cold storage. The difference in molecular weight results in less probability for
25 crystallization. Preferably the molecular weight difference is between 5000 and 20,000 Dalton, most preferably it is about 10,000 Dalton.

In another embodiment recombinant gelatin-like recombinant or synthetic polypeptides of the invention are free from helical structure. This is achieved by allowing only
30 partial or preferably no hydroxylation of the proline residues. Partial hydroxylation means that less than 10% of the prolines are hydroxylated, preferably less than 5%. The absence of helical structure prevents gelling of the gelatin-like polypeptides, even at

low temperatures. This is advantageous in for example vaccine formulations which are dissolved in water before injection. The dissolved vaccine can now be used without the necessity to heat it to prevent gelling.

5 Non gelling gelatin-like polypeptides are also advantageously used in the freeze drying process. In freeze drying of gelatin, the solution is first frozen before the actual freeze drying is started. This process is described in for example US 3,892,876. It is important that the gelatin is frozen in the sol state and not in the gel-state, because otherwise the lyophilized gelatin will not dissolve again after freeze drying. Recombinant gelatin-like
10 proteins of the invention make it possible to freeze dry more concentrated gelatin solutions, resulting in a higher amount of vaccine in the same time, a 10-20% shorter freeze drying time, reducing damage to the physiologically active substance or the gelatin during freeze drying and reducing freeze drying costs.

15 The starting point for the gelatin-like protein for use in the invention can also be an isolated gene encoding a naturally occurring gelatin molecule, which is processed further by recombinant means. Preferably the gelatin-like protein used according to the invention resembles a human native amino acid sequence with this difference that in essence hydroxyproline residues are absent.

20 When produced by recombinant means, especially by expression of recombinant genes in yeasts, the proteins for use according to the invention preferably do not contain a combination of methionine and arginine in 1-4 position (Met-Xay-Xaz-Arg), as such a sequence is sensitive to enzymatic proteolysis.

25 It may be noted that the proteins for use according to the invention can also be partly or wholly produced by methods other than DNA expression, e.g. by chemical protein synthesis.

30 In order to obtain the composition of the invention one or more gelatin-like proteins of the invention are mixed with the physiologically active compound. As an aid in vitrification a saccharide can be added. Preferably this is a disaccharide like sucrose.

Depending on the application also a variety of other compounds can be added like amino acids, other proteins than gelatin, etc.

The composition of the invention comprises an amount of gelatin-like proteins which usually lies in the range from 2-60 weight %.

DESCRIPTION OF THE FIGURES

Fig. 1: Tg of a moving average of n=18 for human COL1A1

Fig. 2: Tg of a moving average of n=27 for human COL1A1

Fig. 3: Tg of a moving average of n=54 for human COL1A1

Fig. 4: Tg of a moving average of n=81 for human COL1A1

Fig. 5: Tg of a moving average of n=54 for human COL1A1; correlation with alanine content

Fig. 6: Tg of a moving average of n=54 for human COL1A1; correlation with leucine + isoleucine content

EXAMPLES

Example 1: Recombinant gelatin-like peptide

A gelatin with an increased glass transition temperature was produced by starting with the nucleic acid sequence that encodes for a part of the gelatin amino acid sequence of human COL1A1-1. The methods as disclosed in EP-A-0926543, EP-A-1014176 and WO01/34646 were used. The sequence of this gelatin according to the invention is given below (SEQ ID NO: 2):

GDRGETGPAGPPGAPGAPGAPGPVGPAGKSGDRGETGPAGPAGPVGP
AGARGPA (amino acid 1034 to 1087 of SEQ ID NO: 1)

Molecular weight: 4590 Da, isoelectric point pI=6.2

This sequence was selected from the total COL1A1-1 sequence (SEQ ID NO: 1) by the method as described in this invention. A glass transition temperature of 208 degrees Celsius was calculated for this selected sequence. The average glass transition

temperature of total COL1A1-1 (SEQ ID NO: 1) is 163 degrees Celsius. Therefore the calculated gain in glass transition temperature is 45 degrees Celsius.

Example 2: Measurement of glass transition temperature

5 The recombinant gelatin as described in example 1 was mixed with sucrose in a ratio of 60/40 wt% gelatin/sucrose, which is typical for MMR vaccine. An aqueous solution of 10% was made of this mixture. This solution was quickly frozen in liquid nitrogen and subsequently it was freeze dried for 48 hours at -55 degrees Celsius. The freeze dried sample was further dried in a vacuum exsiccator with silicagel.

10

DSC (Differential Scanning Calorimetry) was done using a Perkin Elmer DSC 7 instrument under nitrogen atmosphere (flow 20 ml/min). The applied temperature program was:

- 1 minute hold at 60 degrees Celsius

15 - 60 to 230 degrees Celsius at a heating rate of 5 degrees per minute

The glass transition temperature was determined according to the half Cp extrapolated method.

Residual moisture amounts were determined by TGA (Thermo Gravimetric Analysis) using a Perkin Elmer TGA 7 under nitrogen atmosphere (flow 20 ml/min).

20

The applied temperature program was:

- 25-60 degrees Celsius with a heating rate of 5 degrees per minute

- 1 minute hold at 60 degrees Celsius

60 to 300 degrees Celsius at a heating rate of 5 degrees per minute

25

Residual moisture amount of the dry recombinant gelatin/sucrose mixture was found to be in the range of 1-2 wt%.

The glass transition temperature of the dry recombinant gelatin/sucrose mixture was measured to be 130 degrees.

30 As a reference the glass transition temperature of native COL1A1 in the same mixture with sucrose was found to be 116 degrees.

The measured Tg of the mixture comprising the selected recombinant gelatin was thus 14 degrees Celsius higher than the analogous mixture comprising the (non-selected) native gelatin, showing that selection of gelatin-like peptides with a higher calculated Tg also result in mixtures comprising such peptides having a higher measured Tg.

05. 08. 2003

Claims

(87)

1. A lyophilized composition comprising a physiologically active substance and a
5 stabilizer, characterized in that the stabilizer is a recombinant or synthetic gelatin-
like polypeptide comprising at least one stretch of 10 or more consecutive repeats of
Gly-Xaa-Yaa triplets and in which at least 20% of the amino acids are present in the
form of consecutive Gly-Xaa-Yaa triplets and wherein said recombinant
polypeptide has a calculated glass transition temperature of higher than 180 degrees
10 Celsius.
2. A composition as in claim 1 wherein said recombinant or synthetic gelatin-like
polypeptide has a molecular weight between 3,000 Dalton and 80,000 Dalton
preferably between 5,000 Dalton and 60,000 Dalton and more preferably between
10,000 and 40,000 Dalton.
- 15 3. A composition as in claim 1 wherein said recombinant or synthetic gelatin-like
polypeptide has a molecular weight between 3,000 Dalton and 15,000 Dalton
preferably between 5,000 Dalton and 10,000 Dalton and more preferably between
6,000 and 8,000 Dalton.
4. A composition as in the preceding claims wherein the glass transition temperature
20 of the recombinant or synthetic gelatin-like polypeptide is higher than 190 degrees
Celsius preferably higher than 200 degrees Celsius
5. A composition as in the preceding claims wherein the recombinant or synthetic
gelatin-like polypeptide has a bimodal molecular weight distribution
6. A composition as in the preceding claims wherein the recombinant or synthetic
25 gelatin-like polypeptide is free from helical structure
7. A composition as in the preceding claims wherein the number of hydroxyproline
residues in the recombinant or synthetic gelatin-like polypeptide is less than 5% of
the total number of amino acid residues preferably less than 2%
8. A recombinant or synthetic gelatin-like polypeptide comprising at least one stretch
30 of 10 or more consecutive repeats of Gly-Xaa-Yaa triplets and in which at least
20% of the amino acids are present in the form of consecutive Gly-Xaa-Yaa triplets

and wherein said recombinant gelatin-like polypeptide has a calculated glass transition temperature of higher than 180 degrees Celsius.

9. Process for lyophilizing compositions comprising a physiological active substance and a stabilizer characterized in that the stabilizer is a recombinant or synthetic
5 gelatin-like polypeptide comprising at least one stretch of 10 or more consecutive repeats of Gly-Xaa-Yaa triplets and in which at least 20% of the amino acids are present in the form of consecutive Gly-Xaa-Yaa triplets and less than 5% of the total number of amino acid residues are hydroxyproline residues and wherein said
10 recombinant gelatin-like polypeptide has a calculated glass transition temperature of higher than 180 degrees Celsius.

P209992EP

EPO - DG 1

05. 08. 2003

Abstract

(87)

The invention relates to the use of gelatin-like proteins, or polypeptides, with an increased calculated glass transition temperature as stabilisers in lyophilized biological or pharmaceutical compositions.

5

Fig 1

Human COI1A1, n=18




Fig 2

Human COL1A1, n=27




Fig 3

Human COL1A1, n=54




Fig 4

Human COL1A1, n=81




Fig 5

Human COL1A-1, n=54





Fig 6

Human COL1A1, n=54



SEQUENCE LISTING

<110> Fuji Photo Film B.V.

5 <120> Use of gelatin-like proteins as stabiliser

<130> P209992

EPO - DG 1

<160> 2

10

<170> PatentIn version 3.1

05. 08. 2003

<210> 1

<211> 1464

15

<212> PRT

<213> unknown

(87)

<220>

<223> COL1A1

20

<400> 1

Met Phe Ser Phe Val Asp Leu Arg Leu Leu Leu Leu Leu Ala Ala Thr
 1 5 10 15

25

Ala Leu Leu Thr His Gly Gln Glu Glu Gly Gln Val Glu Gly Gln Asp
 20 25 30

30

Glu Asp Ile Pro Pro Ile Thr Cys Val Gln Asn Gly Leu Arg Tyr His
 35 40 45

35

Asp Arg Asp Val Trp Lys Pro Glu Pro Cys Arg Ile Cys Val Cys Asp
 50 55 60

40

Asn Gly Lys Val Leu Cys Asp Asp Val Ile Cys Asp Glu Thr Lys Asn
 65 70 75 80

45

Cys Pro Gly Ala Glu Val Pro Glu Gly Glu Cys Cys Pro Val Cys Pro
 85 90 95

Asp Gly Ser Glu Ser Pro Thr Asp Gln Glu Thr Thr Gly Val Glu Gly
 100 105 110

50

Pro Lys Gly Asp Thr Gly Pro Arg Gly Pro Arg Gly Pro Ala Gly Pro
 115 120 125

55

Pro Gly Arg Asp Gly Ile Pro Gly Gln Pro Gly Leu Pro Gly Pro Pro
 130 135 140

60

Gly Pro Pro Gly Pro Pro Gly Pro Pro Gly Leu Gly Gly Asn Phe Ala
 145 150 155 160

65

Pro Gln Leu Ser Tyr Gly Tyr Asp Glu Lys Ser Thr Gly Gly Ile Ser
 165 170 175

P209992EP

19

Val Pro Gly Pro Met Gly Pro Ser Gly Pro Arg Gly Leu Pro Gly Pro
180 185 190

5 Pro Gly Ala Pro Gly Pro Gln Gly Phe Gln Gly Pro Pro Gly Glu Pro
195 200 205

10 Gly Glu Pro Gly Ala Ser Gly Pro Met Gly Pro Arg Gly Pro Pro Gly
210 215 220

15 Pro Pro Gly Lys Asn Gly Asp Asp Gly Glu Ala Gly Lys Pro Gly Arg
225 230 235 240

20 Pro Gly Glu Arg Gly Pro Pro Gly Pro Gln Gly Ala Arg Gly Leu Pro
245 250 255

25 Gly Thr Ala Gly Leu Pro Gly Met Lys Gly His Arg Gly Phe Ser Gly
260 265 270

30 Leu Asp Gly Ala Lys Gly Asp Ala Gly Pro Ala Gly Pro Lys Gly Glu
275 280 285

35 Pro Gly Ser Pro Gly Glu Asn Gly Ala Pro Gly Gln Met Gly Pro Arg
290 295 300

40 Gly Leu Pro Gly Glu Arg Gly Arg Pro Gly Ala Pro Gly Pro Ala Gly
305 310 315 320

45 Ala Arg Gly Asn Asp Gly Ala Thr Gly Ala Ala Gly Pro Pro Gly Pro
325 330 335

50 Thr Gly Pro Ala Gly Pro Pro Gly Phe Pro Gly Ala Val Gly Ala Lys
340 345 350

55 Gly Glu Ala Gly Pro Gln Gly Pro Arg Gly Ser Glu Gly Pro Gln Gly
355 360 365

60 Val Arg Gly Glu Pro Gly Pro Pro Gly Pro Ala Gly Ala Ala Gly Pro
370 375 380

65 Ala Gly Asn Pro Gly Ala Asp Gly Gln Pro Gly Ala Lys Gly Ala Asn
385 390 395 400

Gly Ala Pro Gly Ile Ala Gly Ala Pro Gly Phe Pro Gly Ala Arg Gly
405 410 415

Pro Ser Gly Pro Gln Gly Pro Gly Gly Pro Pro Gly Pro Lys Gly Asn
420 425 430

P209992EP

20

Ser Gly Glu Pro Gly Ala Pro Gly Ser Lys Gly Asp Thr Gly Ala Lys
435 440 445

5 Gly Glu Pro Gly Pro Val Gly Val Gln Gly Pro Pro Gly Pro Ala Gly
450 455 460

10 Glu Glu Gly Lys Arg Gly Ala Arg Gly Glu Pro Gly Pro Thr Gly Leu
465 470 475 480

15 Pro Gly Pro Pro Gly Glu Arg Gly Gly Pro Gly Ser Arg Gly Phe Pro
485 490 495

Gly Ala Asp Gly Val Ala Gly Pro Lys Gly Pro Ala Gly Glu Arg Gly
500 505 510

20 Ser Pro Gly Pro Ala Gly Pro Lys Gly Ser Pro Gly Glu Ala Gly Arg
515 520 525

25 Pro Gly Glu Ala Gly Leu Pro Gly Ala Lys Gly Leu Thr Gly Ser Pro
530 535 540

30 Gly Ser Pro Gly Pro Asp Gly Lys Thr Gly Pro Pro Gly Pro Ala Gly
545 550 555 560

35 Gln Asp Gly Arg Pro Gly Pro Pro Gly Pro Pro Gly Ala Arg Gly Gln
565 570 575

Ala Gly Val Met Gly Phe Pro Gly Pro Lys Gly Ala Ala Gly Glu Pro
580 585 590

40 Gly Lys Ala Gly Glu Arg Gly Val Pro Gly Pro Pro Gly Ala Val Gly
595 600 605

45 Pro Ala Gly Lys Asp Gly Glu Ala Gly Ala Gln Gly Pro Pro Gly Pro
610 615 620

50 Ala Gly Pro Ala Gly Glu Arg Gly Glu Gln Gly Pro Ala Gly Ser Pro
625 630 635 640

55 Gly Phe Gln Gly Leu Pro Gly Pro Ala Gly Pro Pro Gly Glu Ala Gly
645 650 655

Lys Pro Gly Glu Gln Gly Val Pro Gly Asp Leu Gly Ala Pro Gly Pro
660 665 670

60 Ser Gly Ala Arg Gly Glu Arg Gly Phe Pro Gly Glu Arg Gly Val Gln
675 680 685

65

P209992EP

21

| | | | | | | | | | | | | | | | | | |
|----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|--|
| | Gly | Pro | Pro | Gly | Pro | Ala | Gly | Pro | Arg | Gly | Ala | Asn | Gly | Ala | Pro | Gly | |
| | 690 | | | | | | 695 | | | | | 700 | | | | | |
| 5 | Asn | Asp | Gly | Ala | Lys | Gly | Asp | Ala | Gly | Ala | Pro | Gly | Ala | Pro | Gly | Ser | |
| | 705 | | | | | 710 | | | | | 715 | | | | | 720 | |
| 10 | Gln | Gly | Ala | Pro | Gly | Leu | Gln | Gly | Met | Pro | Gly | Glu | Arg | Gly | Ala | Ala | |
| | | | | | 725 | | | | | 730 | | | | | | 735 | |
| 15 | Gly | Leu | Pro | Gly | Pro | Lys | Gly | Asp | Arg | Gly | Asp | Ala | Gly | Pro | Lys | Gly | |
| | | | | 740 | | | | | 745 | | | | | 750 | | | |
| 20 | Ala | Asp | Gly | Ser | Pro | Gly | Lys | Asp | Gly | Val | Arg | Gly | Leu | Thr | Gly | Pro | |
| | | | 755 | | | | | 760 | | | | | 765 | | | | |
| 25 | Ile | Gly | Pro | Pro | Gly | Pro | Ala | Gly | Ala | Pro | Gly | Asp | Lys | Gly | Glu | Ser | |
| | 770 | | | | | | 775 | | | | | 780 | | | | | |
| 30 | Gly | Pro | Ser | Gly | Pro | Ala | Gly | Pro | Thr | Gly | Ala | Arg | Gly | Ala | Pro | Gly | |
| | 785 | | | | | 790 | | | | | 795 | | | | | 800 | |
| 35 | Asp | Arg | Gly | Glu | Pro | Gly | Pro | Pro | Gly | Pro | Ala | Gly | Phe | Ala | Gly | Pro | |
| | | | | | 805 | | | | | 810 | | | | | 815 | | |
| 40 | Pro | Gly | Ala | Asp | Gly | Gln | Pro | Gly | Ala | Lys | Gly | Glu | Pro | Gly | Asp | Ala | |
| | | | | 820 | | | | | 825 | | | | | 830 | | | |
| 45 | Gly | Ala | Lys | Gly | Asp | Ala | Gly | Pro | Pro | Gly | Pro | Ala | Gly | Pro | Ala | Gly | |
| | | | 835 | | | | | 840 | | | | | 845 | | | | |
| 50 | Pro | Pro | Gly | Pro | Ile | Gly | Asn | Val | Gly | Ala | Pro | Gly | Ala | Lys | Gly | Ala | |
| | | 850 | | | | | 855 | | | | | 860 | | | | | |
| 55 | Arg | Gly | Ser | Ala | Gly | Pro | Pro | Gly | Ala | Thr | Gly | Phe | Pro | Gly | Ala | Ala | |
| | 865 | | | | | 870 | | | | | 875 | | | | | 880 | |
| 60 | Gly | Arg | Val | Gly | Pro | Pro | Gly | Pro | Ser | Gly | Asn | Ala | Gly | Pro | Pro | Gly | |
| | | | | | 885 | | | | | 890 | | | | | 895 | | |
| 65 | Pro | Pro | Gly | Pro | Ala | Gly | Lys | Glu | Gly | Gly | Lys | Gly | Pro | Arg | Gly | Glu | |
| | | | | 900 | | | | | 905 | | | | | 910 | | | |
| 70 | Thr | Gly | Pro | Ala | Gly | Arg | Pro | Gly | Glu | Val | Gly | Pro | Pro | Gly | Pro | Pro | |
| | | | 915 | | | | | 920 | | | | | 925 | | | | |
| 75 | Gly | Pro | Ala | Gly | Glu | Lys | Gly | Ser | Pro | Gly | Ala | Asp | Gly | Pro | Ala | Gly | |
| | 930 | | | | | | 935 | | | | | 940 | | | | | |

Ala Pro Gly Thr Pro Gly Pro Gln Gly Ile Ala Gly Gln Arg Gly Val
 945 950 955 960

5 Val Gly Leu Pro Gly Gln Arg Gly Glu Arg Gly Phe Pro Gly Leu Pro
 965 970 975

10 Gly Pro Ser Gly Glu Pro Gly Lys Gln Gly Pro Ser Gly Ala Ser Gly
 980 985 990

15 Glu Arg Gly Pro Pro Gly Pro Met Gly Pro Pro Gly Leu Ala Gly Pro
 995 1000 1005

20 Pro Gly Glu Ser Gly Arg Glu Gly Ala Pro Gly Ala Glu Gly Ser
 1010 1015 1020

25 Pro Gly Arg Asp Gly Ser Pro Gly Ala Lys Gly Asp Arg Gly Glu
 1025 1030 1035

30 Thr Gly Pro Ala Gly Pro Pro Gly Ala Pro Gly Ala Pro Gly Ala
 1040 1045 1050

35 Pro Gly Pro Val Gly Pro Ala Gly Lys Ser Gly Asp Arg Gly Glu
 1055 1060 1065

Thr Gly Pro Ala Gly Pro Ala Gly Pro Val Gly Pro Ala Gly Ala
 1070 1075 1080

40 Arg Gly Pro Ala Gly Pro Gln Gly Pro Arg Gly Asp Lys Gly Glu
 1085 1090 1095

45 Thr Gly Glu Gln Gly Asp Arg Gly Ile Lys Gly His Arg Gly Phe
 1100 1105 1110

50 Ser Gly Leu Gln Gly Pro Pro Gly Pro Pro Gly Ser Pro Gly Glu
 1115 1120 1125

Gln Gly Pro Ser Gly Ala Ser Gly Pro Ala Gly Pro Arg Gly Pro
 1130 1135 1140

55 Pro Gly Ser Ala Gly Ala Pro Gly Lys Asp Gly Leu Asn Gly Leu
 1145 1150 1155

60 Pro Gly Pro Ile Gly Pro Pro Gly Pro Arg Gly Arg Thr Gly Asp
 1160 1165 1170

65 Ala Gly Pro Val Gly Pro Pro Gly Pro Pro Gly Pro Pro Gly Pro
 1175 1180 1185

| | | | | | | | | | | | | | | | |
|----|-----|------|-----|-----|-----|-----|------|-----|-----|-----|-----|------|-----|-----|-----|
| | Pro | Gly | Pro | Pro | Ser | Ala | Gly | Phe | Asp | Phe | Ser | Phe | Leu | Pro | Gln |
| | | 1190 | | | | | 1195 | | | | | 1200 | | | |
| 5 | Pro | Pro | Gln | Glu | Lys | Ala | His | Asp | Gly | Gly | Arg | Tyr | Tyr | Arg | Ala |
| | | 1205 | | | | | 1210 | | | | | 1215 | | | |
| 10 | Asp | Asp | Ala | Asn | Val | Val | Arg | Asp | Arg | Asp | Leu | Glu | Val | Asp | Thr |
| | | 1220 | | | | | 1225 | | | | | 1230 | | | |
| 15 | Thr | Leu | Lys | Ser | Leu | Ser | Gln | Gln | Ile | Glu | Asn | Ile | Arg | Ser | Pro |
| | | 1235 | | | | | 1240 | | | | | 1245 | | | |
| 20 | Glu | Gly | Ser | Arg | Lys | Asn | Pro | Ala | Arg | Thr | Cys | Arg | Asp | Leu | Lys |
| | | 1250 | | | | | 1255 | | | | | 1260 | | | |
| 25 | Met | Cys | His | Ser | Asp | Trp | Lys | Ser | Gly | Glu | Tyr | Trp | Ile | Asp | Pro |
| | | 1265 | | | | | 1270 | | | | | 1275 | | | |
| 30 | Asn | Gln | Gly | Cys | Asn | Leu | Asp | Ala | Ile | Lys | Val | Phe | Cys | Asn | Met |
| | | 1280 | | | | | 1285 | | | | | 1290 | | | |
| 35 | Glu | Thr | Gly | Glu | Thr | Cys | Val | Tyr | Pro | Thr | Gln | Pro | Ser | Val | Ala |
| | | 1295 | | | | | 1300 | | | | | 1305 | | | |
| 40 | Gln | Lys | Asn | Trp | Tyr | Ile | Ser | Lys | Asn | Pro | Lys | Asp | Lys | Arg | His |
| | | 1310 | | | | | 1315 | | | | | 1320 | | | |
| 45 | Val | Trp | Phe | Gly | Glu | Ser | Met | Thr | Asp | Gly | Phe | Gln | Phe | Glu | Tyr |
| | | 1325 | | | | | 1330 | | | | | 1335 | | | |
| 50 | Gly | Gly | Gln | Gly | Ser | Asp | Pro | Ala | Asp | Val | Ala | Ile | Gln | Leu | Thr |
| | | 1340 | | | | | 1345 | | | | | 1350 | | | |
| 55 | Phe | Leu | Arg | Leu | Met | Ser | Thr | Glu | Ala | Ser | Gln | Asn | Ile | Thr | Tyr |
| | | 1355 | | | | | 1360 | | | | | 1365 | | | |
| 60 | His | Cys | Lys | Asn | Ser | Val | Ala | Tyr | Met | Asp | Gln | Gln | Thr | Gly | Asn |
| | | 1370 | | | | | 1375 | | | | | 1380 | | | |
| 65 | Leu | Lys | Lys | Ala | Leu | Leu | Leu | Lys | Gly | Ser | Asn | Glu | Ile | Glu | Ile |
| | | 1385 | | | | | 1390 | | | | | 1395 | | | |
| 70 | Arg | Ala | Glu | Gly | Asn | Ser | Arg | Phe | Thr | Tyr | Ser | Val | Thr | Val | Asp |
| | | 1400 | | | | | 1405 | | | | | 1410 | | | |
| 75 | Gly | Cys | Thr | Ser | His | Thr | Gly | Ala | Trp | Gly | Lys | Thr | Val | Ile | Glu |
| | | 1415 | | | | | 1420 | | | | | 1425 | | | |

P209992EP

24

Tyr Lys Thr Thr Lys Thr Ser Arg Leu Pro Ile Ile Asp Val Ala
1430 1435 1440

5 Pro Leu Asp Val Gly Ala Pro Asp Gln Glu Phe Gly Phe Asp Val
1445 1450 1455

10 Gly Pro Val Cys Phe Leu
1460

<210> 2

<211> 54

15 <212> PRT

<213> unknown

<220>

20 <223> selected gelatin-like peptide with high Tg

<400> 2

25 Gly Asp Arg Gly Glu Thr Gly Pro Ala Gly Pro Pro Gly Ala Pro Gly
1 5 10 15

Ala Pro Gly Ala Pro Gly Pro Val Gly Pro Ala Gly Lys Ser Gly Asp
20 25 30

30 Arg Gly Glu Thr Gly Pro Ala Gly Pro Ala Gly Pro Val Gly Pro Ala
35 40 45

35 Gly Ala Arg Gly Pro Ala
50

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☒ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☒ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.